

# **Postoperative Instructions**

# **Arthroscopic Partial Meniscectomy**

# Dressings

- Dressings will be removed during your first physical therapy visit.
- Do NOT touch, remove, change or clean any Steri-Strips you may be wearing.
- Do NOT apply or use any ointments such as betadine, Neosporin or peroxide on the wound site.
- Apply additional bandages only if drainage occurs.

### **Bathing**

- Once incisions are dry and no longer draining, you may get them wet in the shower.
  Do NOT get in bath water or a pool until sutures have been removed for at least one week, unless specifically instructed to do so.
- Avoid hot tubs, saunas/steam rooms and hot showers for 4 weeks-they can increase your swelling.

## **Stitches**

- Visible stitches will be removed or clipped at your 2-week follow-up visit in clinic or in PT.
- If you do not have visible sutures to remove, rest assured that they will dissolve under the skin.

#### lce

- Ice your wound site for 20 minutes at a time, 4-5 times a day.
- NEVER place ice directly on your skin. Insulate the site with a towel. Icing helps decrease swelling and providers some pain relief.

#### Swelling

- It is normal to see swelling and/or bruising around your incision for the first 2 weeks.
- Redness, heightened pain, increased temperature (101F or above) and/or abnormal drainage are NOT normal. Please call your doctor's office at once if you develop these conditions.





# **Physical Therapy**

 You will have an appointment scheduled with Physical Therapy within 1 week of surgery.

#### **Medications**

- Please follow all instructions on the bottle.
- Do not drink alcohol while taking pain medication.
- Do not take additional pain medications unless instructed to do so.
- Do not drive while taking pain medication.
- If you have questions about your medications or if you need refills, please reach out to our office at 931-552-4340 and ask to speak to the nurse.

# **Specific Instructions for Knee Care**

# Weight-bearing instructions

Weight-bearing as tolerated to your operative extremity

#### Elevation

Elevate your leg several times a day and at night to decrease swelling.

#### **Immobilization**

• If you were placed in a brace after surgery please wear it until your doctor instructs you to discontinue use. This should also be worn at night.

#### **Contact Numbers**

Clarksville TOA Office: 931-552-4340

#### Additional Instructions:

