



Reverse Shoulder Arthroplasty -WITH SUBSCAPULARIS REPAIR -

General:

- Slingshot for 3 weeks. Regular sling for an additional 3
- ROM should be progressed pain-free; do not push IR or cross-body adduction
- No passive over-pressure

Phase 0 – Quiet

Week 0-2:

- Quiet in sling with elbow/wrist/hand
- Begin active scapular retraction exercises with therapist cueing
- No protraction until 6 weeks

Phase 1 – Passive

Week 2-4:

- Supine External Rotation – 0 degrees or 20 degrees (see orders)
- Supine Forward Elevation – 90 degrees
- No Internal Rotation

Phase 2 – Active

Week 5-6:

- *Active Range of Motion with Passive Stretch to prescribe limits*
- Supine -> Seated Forward Elevation – Full
- Supine -> Seated External Rotation - gradually increase to 45°
- Internal Rotation – gradually increase to full

LUCAS G. TESKE, M.D.

141 Hillcrest Drive | Clarksville, TN 37043 | Phone: 931.552.4340 | Fax : 931.552.0999

TOA

TENNESSEE ORTHOPAEDIC ALLIANCE

Phase 3 – Resisted

Week 7-11:

- *No pendulums and continue with Phase 2*
- *Scapular mobilization*
- External and Internal Rotation – standing forward punch
- Seated Rows/Shoulder Shrugs
- Bicep Curls/Bear Hugs
- Concentrate on deltoid strengthening

Week 12: Weight Training

- Keep hands within eyesight, keep elbows bent
- Minimize overhead activities
- (No military press, pull-down behind head, or wide grip bench)

Return to Activity

Computer	4 weeks
Golf	3 months
Tennis	4 months

LUCAS G. TESKE, M.D.

141 Hillcrest Drive | Clarksville, TN 37043 | Phone: 931.552.4340 | Fax : 931.552.0999

www.TOA.com   